

WITH RELAPSING MS YOUR GOAL IS

TO FIND A WAY TO KEEP

Whether you're new to treatment or ready for a change,

*Clinical Trial 1 (TEMPO) included 1,088 people and Clinical Trial 2 (TOWER) included 1,165 people over 2 years. AUBAGIO 14 mg and 7 mg achieved a significant relative reduction in relapse rate in TEMPO (31%, 31%) and TOWER (36%, 22%) versus placebo. In TEMPO, AUBAGIO 14 mg, AUBAGIO 7 mg, and placebo, the percentage of people who remained free of disability progression were 80%, 78%, and 73% respectively. AUBAGIO 7 mg did not achieve a statistically significant reduction in risk of sustained disability progression. The most common side effects include: headache (16%, 18%, 15%), diarrhea (14%, 13%, 8%), nausea (11%, 8%, 7%), hair thinning or loss (13%, 10%, 5%), and abnormal liver test results (15%, 13%, 9%) for 14 mg, 7 mg, and placebo, respectively.

SEE WHAT ONE PILL  ONCE A DAY CAN DO

INDICATION

AUBAGIO® (teriflunomide) is a prescription medicine used to treat relapsing forms of multiple sclerosis (MS), to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults.

IMPORTANT SAFETY INFORMATION

DO NOT TAKE AUBAGIO IF YOU:

- **Have severe liver problems. AUBAGIO may cause serious liver problems, which can be life-threatening.** Your risk may be higher if you take other medicines that affect your liver. Your healthcare provider should do blood tests to check your liver within 6 months before you start AUBAGIO and monthly for 6 months after starting AUBAGIO. Tell your healthcare provider right away if you develop any of these symptoms of liver problems: nausea, vomiting, stomach pain, loss of appetite, tiredness, yellowing of your skin or whites of your eyes, or dark urine.

Do not take AUBAGIO if you have severe liver problems, are pregnant or of childbearing potential and not using effective birth control, have had an allergic reaction to AUBAGIO or leflunomide, or are taking a medicine called leflunomide for rheumatoid arthritis.

Your healthcare provider will run certain tests before you start treatment. Once on AUBAGIO, your healthcare provider will monitor your liver enzyme levels monthly for the first 6 months.

- **Are pregnant. AUBAGIO may harm an unborn baby.** You should have a pregnancy test before starting AUBAGIO. After stopping AUBAGIO, continue to use effective birth control until you have made sure your blood levels of AUBAGIO are lowered. If you become pregnant while taking AUBAGIO or within 2 years after stopping, tell your healthcare provider right away and enroll in the AUBAGIO Pregnancy Registry at 1-800-745-4447, option 2.

- **Are of childbearing potential and not using effective birth control.**

It is not known if AUBAGIO passes into breast milk. Your healthcare provider can help you decide if you should take AUBAGIO or breastfeed — you should not do both at the same time.

If you are a man whose partner plans to become pregnant, you should stop taking AUBAGIO and talk with your healthcare provider about reducing the levels of AUBAGIO in your blood. If your partner does not plan to become pregnant, use effective birth control while taking AUBAGIO.

SANOFI GENZYME 

ADVERTISEMENT

MOVING FORWARD*

talk to your doctor about once-daily AUBAGIO

Patient Portrayal



Visit myAUBAGIO.com or scan here to learn more

- **Have had an allergic reaction to AUBAGIO or a medicine called leflunomide.**
- **Take a medicine called leflunomide for rheumatoid arthritis.**

AUBAGIO may stay in your blood for up to 2 years after you stop taking it. Your healthcare provider can prescribe a medicine that can remove AUBAGIO from your blood quickly.

Before taking AUBAGIO, talk with your healthcare provider if you have: liver or kidney problems; a fever or infection, or if you are unable to fight infections; numbness or tingling in your hands or feet that is different from your MS symptoms; diabetes; serious skin problems when taking other medicines; breathing problems; or high blood pressure. Your healthcare provider will check your blood cell count and TB test before you start AUBAGIO. Talk with your healthcare provider if you take or are planning to take other medicines (especially medicines for treating cancer or controlling your immune system), vitamins or herbal supplements.

AUBAGIO may cause serious side effects, including: reduced white blood cell count — this may cause you to have more infections; numbness or tingling in your hands or feet that is different from your MS symptoms; allergic reactions, including serious skin problems;

breathing problems (new or worsening); and high blood pressure. Patients with low white blood cell count should not receive certain vaccinations during AUBAGIO treatment and 6 months after.

Tell your doctor if you have any side effect that bothers you or does not go away.

The most common side effects when taking AUBAGIO include: headache; diarrhea; nausea; hair thinning or loss; and abnormal liver test results. These are not all the side effects of AUBAGIO. Tell your healthcare provider about any side effect that bothers you.

Consult your healthcare provider if you have questions about your health or any medications you may be taking, including AUBAGIO.

You may report side effects to the FDA at 1-800-FDA-1088.

Please see Medication Guide for AUBAGIO on adjacent pages and full Prescribing Information, including boxed WARNING, available on www.aubagio.com.



PILL ACTUAL SIZE

Once-daily
AUBAGIO®
(teriflunomide) 14mg tablets

AUBAGIO is available in 14 mg and 7 mg tablets.

©2020 Genzyme Corporation. All rights reserved. AUBAGIO, Sanofi and Genzyme registered in U.S. Patent and Trademark Office. SAUS.AUBA.20.01.0288. March 2020.